

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Medical Microbiology and Tropical Diseases

Journal homepage: <https://www.ijmmt.org/>

## Original Research Article

## Biofilm production and antibiotic resistance pattern among bacterial isolates from wound samples in a rural tertiary care teaching hospital

Venkatesha D<sup>1</sup>, Dhanalakshmi T. A<sup>1</sup>, Shakthi R<sup>1,\*</sup><sup>1</sup>Dept. of Microbiology, Adichunchanagiri Institute of Medical Sciences, Mandya, Karnataka, India

## ARTICLE INFO

## Article history:

Received 29-06-2021

Accepted 02-08-2021

Available online 01-09-2021

## Keywords:

Antibiotic resistance

Biofilm

Congo red agar

Wound infection

## ABSTRACT

**Introduction:** Pyogenic wound infections are the one of the leading cause of morbidity and mortality worldwide. Some of the common etiological agents responsible are *Staphylococcus aureus*, *Streptococcus pyogenes*, *Escherichia coli*, *Klebsiella app.*, *Proteus spp.*, *Pseudomonas spp.*, and *Acinetobacter spp.* The antimicrobial resistance has become a global challenge and the resistant pathogen poses a grave threat to the public health worldwide. Pyogenic bacteria producing biofilm has a potential to cause significant mortality and morbidity in human.

**Aim:** The present study was carried out to determine the bacteriological spectrum of wound infections and their antibiogram to commonly used antibiotics and to detect the biofilm production by the isolates.

**Materials and Methods:** This cross sectional study was carried out in the department of Microbiology, Adichunchanagiri institute of Medical sciences from September 2016 to August 2017. Two hundred and forty samples from various wounds were collected and processed as per standard procedures and biofilm production was detected by Congo red agar method.

**Results :** Out of 240 pus isolates, *Staphylococcus* species were the most commonly isolated (48.85%) followed by *Pseudomonas* species (11.7%). Biofilm was produced by 49.2% isolates. Majority of Gram negative bacilli were susceptible to Colistin (100%) followed by Tigecycline (Biofilm producers 75%, biofilm non producers 66.7%). All Gram positive isolates were susceptible to Vancomycin and Teicoplanin (100%) followed by Linezolid (biofilm producer 98.8%, biofilm non producer 97.8%).

**Conclusion:** Routine surveillance for wound infections along with early identification and adopting efficient control protocol against biofilm forming organism plays an important role in the prevention of the most serious infections.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Pyogenic wound infections are the one of the leading cause of morbidity and mortality worldwide.<sup>1,2</sup> Wound sepsis rate in India is around 10-33%. Infections of the wound also contribute huge economic burden because of prolonged hospital stay.<sup>3</sup> Upsurge of antimicrobial resistance has provided a new angle to the current problem of wound infections.<sup>4</sup> Biofilms have enormous negative impact on

health care system.<sup>5</sup> Formation by biofilms by wound isolates impair healing of the wound, reduce host immune response and further add on to the development of antibiotic resistance as biofilms impede delivery of antibiotics, cause degradation of antibiotics, promote horizontal transfer of resistance genes<sup>1</sup>. Biofilms complicate patient treatment.<sup>6</sup> Data on biofilm formation by isolates is required to modify treatment modality and the outcome of clinical condition.<sup>7</sup> The spectrum of bacteria causing wound infections and their antibiotic susceptibility patterns exhibit geographic variability and also changing trends noted with respect to

\* Corresponding author.

E-mail address: [drshakthir@bgsaims.edu.in](mailto:drshakthir@bgsaims.edu.in) (Shakthi R).

time.<sup>8,9</sup> Updated information regarding the bacteriological profile and their antibiogram is valuable for implementing strategies towards empiric treatment of wound infections, in adopting efficient control protocols and in formulation of suitable antibiotic policies for treatment of wound infections in the region.<sup>6,8</sup> With this background the present study was carried out to determine the bacteriological spectrum of wound infections and their antibiogram to commonly used antibiotics and to detect the biofilm production by the isolates.

## 2. Materials and Methods

The present cross sectional study was carried out in the department of Microbiology, Adichunchanagiri institute of Medical sciences from September 2016 to August 2017. Two hundred and forty pus samples from various wounds were collected and submitted to microbiology laboratory of the hospital and were processed as per standard procedures.<sup>10</sup> Both inpatients and outpatients were included in the study.

Repeated isolates from the same patient and patients who were on antibiotic therapy or had history of antibiotic intake within one week prior to sample collection and anaerobic isolates were excluded from the study. Informed consent from patients and ethical clearance from the institution were obtained for the study.

Specimens were inoculated on to Blood agar and MacConkey agar plates (procured from HiMedia Mumbai India) and incubated for 48 hours at 37<sup>0</sup>C under aerobic conditions. Isolates were identified by standard microbiological methods.<sup>10</sup> Antibiotic susceptibility testing was carried out by Kirby Bauer disc diffusion method as per Clinical Laboratory Standards Institute guidelines with ATCC *Staphylococcus aureus* ATCC 25923 and *Escherichia coli* ATCC 25922 as control strains.<sup>11,12</sup>

For biofilm detection, *S.epidermidis* ATCC 35984 and *S.epidermidis* ATCC 12228 were used as positive and negative controls respectively.<sup>13</sup>

Isolates were tested for biofilm production by Congo red agar method. Isolates were inoculated onto Congo red agar plate (procured from HiMedia Mumbai, India) and were incubated for 48 hours at 37<sup>0</sup>C under aerobic conditions. The appearance of black colonies with dry crystalline consistency was taken as positive for slime production. Isolates producing very dark coloured colonies were interpreted as strong biofilm producers. Those bacteria forming black colonies were considered as moderate biofilm producers and those producing almost black colonies were noted as weak biofilm producers. Isolates forming red colonies were considered as non-biofilm producers. Each test was interpreted by two different observers<sup>14,15</sup> (Figure 1).

## 2.1. Statistical analysis

Statistical analysis was done using Microsoft excel. The data analysis involved transcription, preliminary data inspection, content analysis and interpretation. Percentages were used in this study to analyse variables

## 3. Results

Out of 240 pus isolates studied, *Staphylococcus* species were the most commonly isolated bacteria (48.85) followed by *Pseudomonas* species (11.7%) (Table 1).

Biofilm was produced by 49.2% of total isolates. Among biofilm producers 39% were strong biofilm producers followed by mild (37.3%) and moderate biofilm producers (23.7%). Among Gram negative bacilli isolated, biofilm production was observed more among *E. coli* and *Klebsiella* species (29.7% each) followed by *Citrobacter species* (18.9%).(Figure 2).

Out of total Gram positive cocci isolated in the study, majority of biofilm production was seen among Methicillin Resistant *Staphylococcus aureus* (MRSA) isolates (48.1%) followed by Methicillin Resistant Coagulase negative *Staphylococci* (22.2%). (Figure 3)

Majority of Gram negative bacilli were found sensitive to Colistin, Tigecycline, and aminoglycosides followed by Imipenem. (Table 2)


Majority of Gram positive cocci were sensitive to Teicoplanin, Vancomycin and Linezolid (Table 3). Majority of *Pseudomonas* species were sensitive to ciprofloxacin followed by carbapenems and aminoglycosides. (Figure 4)

**Table 1:** Bacterial pathogens isolated from pus specimen


Bacteria isolates	Number (%)
<i>Staphylococci</i> species	117 (48.8)
<i>Pseudomonas</i> species	28 (11.7)
<i>Klebsiella</i> species	24 (10.0)
<i>E.coli</i>	21(8.8)
<i>Citrobacter species</i>	17(7.1)
Gram negative non-fermenters other than <i>Pseudomonas</i> and <i>Acinetobacter</i> spp	10 (4.2)
<i>Enterococcus</i> species	9 (3.8)
<i>Enterobacter</i> species	4 (1.7)
<i>Proteus</i> species	3 (1.3)
<i>Streptococcus</i> species	2 (0.8)
<i>Acinetobacter</i> species	2 (0.8)
<i>Morganella</i> species	2 (0.8)
<i>Providencia</i> species	1 (0.4)
Total	240 (100)

## 4. Discussion


Wound infection is one of the most common and serious complication among the hospital acquired infections. Wound infection can increase the length of hospital stay and accounts for mortality rate up to 70-80%.<sup>12</sup>



**Fig. 1:** Congored agar method: a) Strong biofilm producer; b) Moderate biofilm producer; c) Weak biofilm producer; d) Non biofilm producer



**Fig. 2:** Biofilm forming Gram negative bacilli




**Fig. 3:** Biofilm forming Gram positive cocci

**Table 2:** Sensitivity pattern of Gram negative bacilli (%)

Antibiotics	Biofilm positive	Biofilm Negative
Ampicillin	2.7	1.4
Amoxicillin -Clavulanic acid	2.9	2.8
Piperacillin-Tazobactam	37.8	56.5
Ceftriaxone	35.3	13.8
Ceftazidime	29.7	32.4
Cefepime	44.4	43.1
Imipenem	50	72.6
Ciprofloxacin	38.2	49.3
Ofloxacin	40.6	53.7
Gentamicin	58.3	67.2
Amikacin	67.6	65.3
Tobramycin	63.9	77.5
Tigecycline	75	66.7
Colistin	100	100

**Table 3:** Sensitivity pattern of Gram positive cocci (%)

Antibiotics	Biofilm Positive	Biofilm Negative
Penicillin	6.4	13.6
Amoxicillin-Clavulanic acid	11.8	20.9
Ceftriaxone	0	0
Cefotaxime	17.2	27.9
Erythromycin	22.2	32.6
Clindamycin	45.2	53.5
Ciprofloxacin	13.7	29.5
Ofloxacin	7.4	36.7
Gentamicin	41.7	55.6
Amikacin	50.6	62.2
Tetracycline	78.4	73.3
Chloramphenicol	69.6	71.8
Cotrimoxazole	38.5	43.2
Vancomycin	100	100
Linezolid	98.8	97.8
Teicoplanin	100	100



**Fig. 4:** Sensitivity pattern observed among *Pseudomonas* species

In the present study, a dominance of Gram Positive bacteria as the causative agent of pyogenic lesions was seen similar to studies by Tiwari et al., Lee et al., 2009.<sup>16,17</sup> This is in contrast to other studies where Gram negative bacteria are most common pathogens isolated.<sup>12,18</sup> *Staphylococcus* forms the normal microbial flora of the skin and anterior nares and hence can easily contaminate the wound.<sup>2,5,6</sup>

*Staphylococcus aureus* (48.85%) was the most common Gram positive isolate which similar to the various studies.<sup>12,16,17</sup> The prevalence of MRSA (39%) reported in our study is in accordance with the studies of Kshetry et al.,(37.6%), Sanjana et al.,(37.6%), Dibah et al.,(46.3%).<sup>19–21</sup> However, lower rates were reported by Rozina AK et al., Tiwari et al., Lee et al.,<sup>12,16,17</sup>. The difference in the rates of isolation of MRSA might be due to the difference in the level of irrational antibiotic use, level of hygienic condition maintained in different hospitals and effective implementation of hand hygiene program.<sup>12</sup>

Among Gram Negative bacilli, the most common isolate was *Pseudomonas* species (11.7%) followed by *Klebsiella* species (10%) and *Escherichia coli* (8.8%). This is in accordance with the study of Mukhopadhyay et al.,<sup>22</sup> but is in contrast with the study of Pal K et al., where *Escherichia coli* was the most common Gram negative isolate.<sup>23</sup> This inconsistency in bacterial pattern may be due to regional variation of bacterial profile, habits of the local people and also due to the fact that all the Indian studies done so far involve community set up and were not centered on hospitals.<sup>24</sup>

Antimicrobial resistance is an innate feature of bacterial biofilm that, in addition to the increasing resistance among clinical strains, may complicate patient treatment.<sup>6</sup> In this study biofilm was produced by 49.2% of total isolates. Out of total Gram positive cocci isolated in the study, majority of biofilm production was seen among Methicillin Resistant *Staphylococcus aureus* isolates (48.1%), which is similar to the studies of Shrestha et al (44.9%) and Ansari et al (43.1%).<sup>25,26</sup>

Majority of Gram negative bacilli were found sensitive to Colistin, Tigecycline, and aminoglycosides followed by Imipenem which is similar to various studies.<sup>8,9,27</sup> Majority of *Pseudomonas* species were sensitive to ciprofloxacin followed by carbapenems and aminoglycosides, this is in accordance with Sukumar N, et al.,; Sharma V et al., and Amabegaum B et al.,<sup>8,9,28</sup>

Majority of Gram positive cocci were sensitive to Teicoplanin, Vancomycin and Linezolid which is in agreement with other studies.<sup>1,9,29,30</sup>

The prevalence and pattern of Antimicrobial resistance among wound isolates show variability according to geographic location, endemicity of pathogen in the locality and climate conditions.

## 5. Limitation of the Study

In this study, clinical correlation was not done. Small sample size did not allow us to conduct advanced statistical analysis, which could have potentially strengthened this study

## 6. Conclusion

In the present study, pyogenic wound infections were mainly caused by *S. aureus*, *Pseudomonas* spp, *Klebsiella* spp., and *Escherichia coli*. Both Gram positive and Gram negative bacteria with biofilm production showed modest increase in resistance to few antibiotics compared to those without biofilm production.

Continuous surveillance is necessary to update the knowledge of antimicrobial susceptibility pattern of clinical isolates to provide the appropriate dose regimen and treatment schedule against pyogenic wound infections and to limit the expanding threat of drug resistance. Early identification and adopting efficient control protocol against biofilm forming organism can prevent the most serious nosocomial infections and improve the outcome of the condition.

## 7. Conflict of Interest

The authors declare that there are no conflicts of interest in this paper.

## 8. Source of Funding

None

## References

1. Rijal BP, Satyal D, Parajuli NP. High burden of antimicrobial resistance among bacteria causing pyogenic wound infections at a tertiary care hospital in Kathmandu, Nepal. *J Pathog*. 2017;doi:10.1155/2017/9458218.
2. Kassam NA, Damian DJ, Kajeguka D, Nyombi B. Spectrum and antibiogram of bacteria isolated from patients presenting with infected wounds in a tertiary Hospital, northern Tanzania. *BMC Res Notes*. 2017;10:757. doi:10.1186/s13104-017-3092-9.
3. Sukumar N, Rajesh S. Aerobic bacterial isolates and their antibiotic susceptibility pattern from pus samples in a tertiary care Government Hospital in Tamil Nadu, India. *Int J currMicrobiol App Sci*. 2017;6(6):423–42.
4. Biradar A, Farooqui F, Prakash R, Khaqri S, Itag I. Aerobic bacteriological profile with antibiogram of pus isolates. *Indian J Microbiol Res*. 2016;3(3):245–9. doi:10.5958/23945478.2016.00054.6.
5. Kaur DC, Wankhede. Biofilm formation and antimicrobial susceptibility pattern of Methicillin Resistant *Staphylococcus aureus* from wound infection. *Asian Pc J Health Sci*. 2014;1(4):322–8.
6. Kaur DC, Khare A. Biofilm formation and antibiotic susceptibility pattern in MRSA strains in a tertiary care rural hospital. *Indian J Basic Appl Med Res*. 2013;1(3):37–44.
7. Asati S, Chaudhary U. Prevalence of biofilm producing aerobic bacterial isolates in burn wound infections at a tertiary care hospital in northern India. *Ann Burns Fire Disasters*. 2017;30(1):39–44.
8. Trojan R, Razdan L, Singh N. Antibiotic Susceptibility Patterns of Bacterial isolates from Pus samples in a Tertiary Care Hospital of. *Int J Microbiol*. 2016;doi:10.1155/2016/9302692.

9. Sharma V, Parihar G, Sharma V, Sharma H. A study of various isolates from Pus sample with their antibiogram from Jin Hospital, Ajmer. *J Dent Med Sci*. 2015;14(10):64–8.
10. Collee JG, Mackie TJ, McCartney JE, Simmons A. Mackie & McCartney practical medical microbiology. In: 14th Edn. Edinburgh(UK):Churchill Livingstone; 1999.
11. Clinical and Laboratory Standards. Performance Standards for Antimicrobial Susceptibility testing; Twenty fifth Informational Supplement. CLSI Document M100-S25. Wayne PA, USA: Clinical and Laboratory Standards Institute; 2015.
12. Rozina AK, Mahwish J, Mohammed K. Bacteriological profile and antibiogram of isolates from pus samples in a tertiary care centre. *Int J Curr Microbiol App Sci*. 2018;7(1):387–94.
13. Niveditha S, Pramodhini S, Umadevi S, Kumar S, Stephen S. The isolation and the biofilm formation of uropathogens in the patients with catheter associated urinary tract infections. *J Clin Diagn Res*. 2012;6(9):1478–82.
14. Murugan K, Usha M, Al-Sohaibani A, Chandrasekaran M. Biofilm forming conjunctivitis multi drug resistant Staphylococcus spp among the ocular patients. *Pol J Microbiol*. 2010;59(4):233–9.
15. Ferreira AA, Tette PAC, Mendonca RCS, Soares ASC, Carvalho MM. Detection of exopolysaccharide production and biofilm related genes in Staphylococcus species isolated from poultry processing plant. *Food Sci Technol*. 2014;34(4):710–6.
16. Tiwari HK, Das AK, Sapkota D, Pahwa V, Sivarajan K. Methicillin resistant Staphylococcus aureus: prevalence and antibiogram in a tertiary care hospital in western Nepal. *J Infect Dev Ctries*. 2009;3(9):681–4.
17. Lee CY, Chen PY, Huang FL, Lin F. Microbiologic spectrum and susceptibility pattern of clinical isolates from the pediatric intensive care unit in single medical centre-6years experience. *J Microbiol Immunol Infect*. 2009;42(2):160–5.
18. Zubair M, Malik A, Ahmad J. Clinico-microbiological study and antimicrobial drug resistance profile of diabetic foot infections in North India. *Foot*. 2011;21(1):6–14.
19. Kshetry AO, Pant ND, Bhandarirkhatri S, Shrestha KL, Upadhaya SK. Minimum inhibitory concentration of Vancomycin to methicillin resistant Staphylococcus aureus isolated from different clinical samples at a tertiary care hospital in Nepal. *Antimicrobial Resist Infect Control*. 2016;5:27. doi:10.1186/s13756-016-0126-3.
20. Sanjana R, Shah R, Chaudhary N, Singh Y. Prevalence and antimicrobial susceptibility pattern of Methicillin resistant Staphylococcus aureus(MRSA) in CMS- teaching hospital: a preliminary report. *J Coll Med Sci-Nepal*. 2010;6(1):1–6.
21. Dibah S, Arzanlou M, Jannati E, Shapouri R. Prevalence and antimicrobial resistance pattern of Methicillin resistant Staphylococcus aureus (MRSA) strains isolated from clinical specimens in Ardabil, Iran. *Iranian J Microbiol*. 2014;6(3):163–8.
22. Mukhopadhyay M, Podder S, Bhattacharya S. Microbial contamination of Indian currency notes and coins in Kolkata, West Bengal - A Survey. *Int J Sci Res*. 2015;4(8):618–20.
23. Pal K, Das NS, Bhattacharya S. Bacteriological profile of Indian currency circulating in a tertiary care hospital in rural Bengal. *JRRMS*. 2013;3(2):23–7.
24. Afroz Z. Bacteriological Profile and Antimicrobial Susceptibility Pattern of Indian Currency Circulating in a Tertiary Care Hospital of South India. *Int J Curr Microbiol App Sci*. 2018;7(4):1828–34.
25. Shrestha B, Pokhrek BM, Mohapatra T. Phenotypic characterization of nosocomial isolates of Staphylococcus aureus with reference to MRSA. *J Infect Dev Ctries*. 2009;3(7):554–60.
26. Ansari S, Nepal HP, Gautam R, Rayamajhi N, Shrestha S, Upadhyay G, et al. Threat of drug resistant Staphylococcus aureus to health in Nepal. *BMC Infect Dis*. 2014;14:157. doi:10.1186/1471-2334-14-157.
27. Kassam NA, Damian JD, Debora K, Balthazar N, Gibson S. Spectrum and antibiogram of bacteria isolated from patients presenting with infected wounds in a tertiary hospital, northern Tanzania. *BMC Res Notes*. 2017;10:757. doi:10.1186/s13104-017-3092-9.
28. Amabegaum B, Farooqui F, Ravichandra P, Sayeda YK, Ifran I. Aerobic bacteriological profile with antibiogram of pus isolates. *Indian J Microbiol Res*. 2016;3(3):245–9.
29. Mohammed A, Seid ME, Gebrecherkos T, Tiruneh M, Moges F. Bacterial Isolates and Their Antimicrobial Susceptibility Patterns of Wound Infections among Inpatients and Outpatients Attending the University of Gondar Referral Hospital, Northwest Ethiopia. *Int J Microbiol*. 2017;p. 8953829. doi:10.1155/2017/8953829.
30. Belbase A, Narayan DP, Krishus N, Bihusan N, Rikesh B, Reena B, et al. Antibiotic resistance and biofilm production among the strains of Staphylococcus aureus isolated from pus /wound swab samples in a tertiary care hospital in Nepal. *Ann Clin Microbiol Antimicrob*. 2017;16(1):15. doi:10.1186/s12941-017-0194-0.

### Author biography

**Venkatesha D**, Professor and Head

**Dhanalakshmi T. A**, Professor

**Shakthi R**, Assistant Professor

**Cite this article:** Venkatesha D, Dhanalakshmi T. A, Shakthi R. Biofilm production and antibiotic resistance pattern among bacterial isolates from wound samples in a rural tertiary care teaching hospital. *IP Int J Med Microbiol Trop Dis* 2021;7(3):160-164.