

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Medical Microbiology and Tropical Diseases

Journal homepage: <https://www.ijmmt.org/>

## Original Research Article

## The predictive role of combined adenosine deaminase activity in Serum and body fluids to diagnose extrapulmonary tuberculosis

Sateesh K<sup>1,\*</sup>, Sathya Anandam<sup>2</sup><sup>1</sup>Dept. of Microbiology, Kamineni Institute of Medical Science, Narketpally, Telangana, India<sup>2</sup>Dept. of Microbiology, Karpagam Faculty of Medical sciences and Research, Coimbatore, Tamil Nadu, India

## ARTICLE INFO

## Article history:

Received 17-05-2023

Accepted 26-06-2023

Available online 18-07-2023

## Keywords:

Serum ADA

Body Fluids ADA

Extrapulmonary TB

predictive biomarker

## ABSTRACT

**Background:** EPTB comprises 10-15% of all TB cases in developing countries. Diagnosis of TB from body fluids like pleural, peritoneal and cerebrospinal fluid (CSF) is challenging as all these fluid samples possess very few bacilli.

**Aim:** To determine the role of adenosine deaminase (ADA) assay for reliable prediction of EPTB from different body fluids, particularly in low-resource areas with high disease prevalence.

**Materials and Methods:** This prospective study was out in a rural medical college hospital. An enzymatic deamination method in a kinetic manner was used to monitor the ADA activity. The study processed 100 serum samples from 50 Suspected TB patients and 50 from the control group and 100 samples of body fluids from 50 Suspected TB patients and 50 control samples. Data were recorded in MS Excel sheets, and statistical analysis was performed using MS Office software.

**Results:** Out of 50 serum samples from the suspected TB patient and control groups, 48 (96%) and 17 (34%) were positive for ADA, respectively. Out of 50 samples of body fluids obtained from both suspected TB patients and the control group, 16 (32%) and 3 (06%) were positive for ADA, respectively. Serum ADA positivity was significantly high in suspected TB patients as compared to the control group

**Conclusions:** In our study, observations suggest that serum and serosal fluid Adenosine deaminase (ADA) measurement has good prediction potential for EPTB. Hence, it can be used as a supportive surrogate marker for challenging to diagnose extrapulmonary TB. ADA activity in body fluids is also a sensitive biomarker, especially when combined with serum ADA levels and may become a routine investigation for early detection of extrapulmonary TB.

**Key Messages:** Serum and serosal fluid Adenosine deaminase (ADA) measurements have good prediction potential for PTB & EPTB.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Tuberculosis (TB) is a protean disease caused by *Mycobacterium tuberculosis* (*M. tuberculosis*). Usually, TB patients present with pulmonary manifestations. Nevertheless, extrapulmonary tuberculosis (EPTB) is not uncommon in TB-endemic countries like India and other south-east Asian countries. EPTB comprises 10-15% of

all TB cases in developing countries.<sup>1,2</sup> Diagnosis of TB from body fluids like pleural, peritoneal and cerebrospinal fluid (CSF) is challenging as all these fluid samples possess very few bacilli. Conventional methods like microscopy and culture are widely used for diagnosis, but the sensitivity of AFB (Acid-fast bacilli) smear is only 5-20%, and culture takes about three to six weeks which may prolong the initiation of treatment.<sup>3-6</sup>

\* Corresponding author.

E-mail address: [drsateesh2006@gmail.com](mailto:drsateesh2006@gmail.com) (Sateesh K).

Owing to this fact, newer tests have been developed like antigen-antibody detection, antibody in lymphocyte supernatant (ALS) assay, cellular IFN $\gamma$  release assays (IGRAs) and T-Spot.<sup>7,8</sup> However, the reliability of these tests in diagnosing active TB disease is not proven yet. With the advent of molecular technology, polymerase chain reaction (PCR) has been developed to detect *M. tuberculosis* rapidly. PCR can reliably detect a very low concentration of organisms in extrapulmonary samples.<sup>9–11</sup> However, the requirement of dedicated laboratory areas, rigorous quality control, and the high test cost limits its routine use in resource-poor countries.

In this regard, our study aimed to study the role of adenosine deaminase (ADA) assay for reliable prediction of EPTB from different body fluids, particularly in resource-poor areas and where the disease is prevalent.<sup>12–15</sup>

## 2. Materials and Methods

The study was conducted from October 2018 to November 2019 in the Microbiology department of a tertiary care rural Hospital, utilizing serum samples and body fluids like Pleural fluid, ascitic fluid, CSF, Urine and pericardial fluid received in the microbiology laboratory from all clinically suspected extrapulmonary tuberculosis patients. Serum samples and body fluids like Pleural fluid, ascitic fluid, CSF, Urine and pericardial fluid were collected from randomly selected patients suffering from other confirmed non-tuberculosis diseases like cancer, cirrhosis, post pneumonic effusions, pyogenic meningitis, and peritonitis were used as controls.

### 2.1. Ethical consideration

Ethical clearance from the Institutional Ethical Review Committee was taken and informed consent from the patients was waived by an ethical committee before commencing the study.

### 2.2. Specimen preparation

#### 2.2.1. Serum/ CSF/ Body Fluid

No special preparation of the patient is required prior to sample collection by approved techniques.

ADA is reported to be stable in Serum for three days at 2–8°C and in biological fluids for two days at 2–8°C, as after this, ammonia may be released in the samples even without any microbial contamination. Hence all samples were processed on the same collection day to prevent false negative results.





### 2.3. Estimation of ADA levels

#### 2.3.1. Principle

ADA Assay is based on the enzymatic deamination of adenosine to Inosine. Purine nucleoside phosphorylase

(PNP) enzyme later converts it to Hypoxanthine. Furthermore, Hypoxanthine gets converted to uric acid and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by xanthine oxidase (XOD). In the presence of peroxidase (POD), H<sub>2</sub>O<sub>2</sub> is further reacted with N-Ethyl-N-(2-hydroxy-3-sulfoethyl)-3-methyl aniline (EHSPT) and 4-amino antipyrine (4-AA) to generate quinone dye which is thereby monitored in a kinetic manner.

Below, the entire enzymatic reaction scheme is shown.



One unit of ADA is defined as the amount of ADA that generates one  $\mu\text{mole}$  of Inosine from adenosine per min at 37°C.

#### 2.3.2. Reagent

Micropress ADA-MTB is a reagent for laboratory use only.

ADA-MTB comprises of:

1. L1 - ADA-MTB Reagent - Buffer Reagent, ready to use.
2. L2 - ADA-MTB Reagent - Adenosine Reagent, ready to use.
3. L3 - ADA-MTB Reagent - Phenol Reagent.
4. L4 - ADA-MTB Reagent - Hypochlorite Reagent.
5. L5 - ADA-MTB Standard - ADA Standard, ready to use.

**Specimen preparation:** No special preparation of the patient is required prior to sample collection by approved techniques

ADA is reported to be stable in Serum for three days at 2–8°C and in biological fluids for two days at 2–8°C, as after this, ammonia may be released in the samples even without microbial contamination.

### 2.4. Test procedure

1. Pipette into clean, dry test tubes labelled Blank (B), Standard (S), Sample Blank (SB) and Test (T), as shown in Table 1.
2. Mix well and incubate at 37°C for precisely 60 minutes, and then follow Table 2.
3. Mix well and incubate at 37°C for 15 minutes or at RT for 30 minutes.
4. Measure the absorbance of the Blank (Abs. B), Standard (Abs. S), Sample Blank (Abs. SB) and Test (Abs. T) against distilled water.

**Table 1:** Addition sequence of reagents

| Addition sequence | B (ml) | S (ml) | SB (ml) | T (ml) |
|-------------------|--------|--------|---------|--------|
| Buffer reagent    | 0.20   | 0.20   | -       | -      |
| Adenosine Reagent | -      | -      | 0.20    | 0.20   |
| Deionized water   | 0.02   | -      | -       | -      |
| Standard          | -      | 0.02   | -       | -      |
| Sample            | -      | -      | -       | 0.02   |

**Table 2:** Addition sequence of reagents

|                              | B (ml) | S (ml) | SB (ml) | T (ml) |
|------------------------------|--------|--------|---------|--------|
| Working Phenol Reagent       | 1.00   | 1.00   | 1.00    | 1.00   |
| Sample                       | -      | -      | 0.02    | -      |
| Working Hypochlorite Reagent | 1.00   | 1.00   | 1.00    | 1.00   |

2.5. Calculations

$$\text{Total ADA activity in U/L} = \frac{\text{Abs. T} - \text{Abs. SB} \times 50}{\text{Abs. S} - \text{Abs. B}}$$

**Table 3:** Reference values:

| Sample   | Interpretation | Result         |
|--|----------------|----------------|
| Serum, Plasma, Pleural, Pericardial and Ascitic Fluids | Normal         | <30U/L         |
|  | Suspect        | 30U/L to 40U/L |
|  | Strong Suspect | >40U/L - 60U/L |
| CSF  | Positive       | >60U/L         |
|  | Normal         | <10U/L         |
|  | Positive       | >10U/L         |

2.6. Details of data tabulation and statistical analysis employed

Data was recorded in MS Excel sheets, and statistical analysis was done using MS Office software.

3. Results


We included 50 suspected EPTB patient’s Serum samples and body fluids as cases and compared them with 50 randomly selected patient’s Serum samples and body fluids suffering from other confirmed non-tuberculosis diseases

The age wise distribution of 100 samples was 51-60yrs (27%), 31-40yrs(20%), 41-50yrs (18%), 20-30yrs (13%) and 61-70yrs (13%) and 71-90yrs (9%). Concerning gender, 75% of the patients in this study were males and 25% were females.

ADA activity in both cases and control was compared, and it was found that 48(96%) serum samples and 16 (32%) body fluid samples obtained from the suspected EPTB patient group were positive for ADA. In comparison, 17 (34%) serum samples and 3(06%) body fluid samples in the control group showed ADA activity.


Out of 50 serum samples in the suspected EPTB patient group, 48 (96%) were positive, and out of 50 serum samples in the control group, 17 (34%) were positive for ADA enzyme. Out of 50 body fluid samples in the suspected EPTB patient group, 16 (32%) were positive, and out of 50 body fluids in the control group, 3 (06%) were positive for ADA enzyme.

In the suspected TB patient group, combined ADA activity in both Serum and body fluid samples was analyzed. It was found that 14 (28%) were positive for ADA in the same patient’s serum and body fluids. However, 34(68%) of the sample were positive in Serum and no ADA activity in body fluids. No samples were negative for ADA activity in both Serum and body fluid. In the Control group, 30 (60%) samples were negative for ADA activity in both Serum and body fluid. However, 17(34%) samples were positive for only Serum ADA. 3(6%) samples showed ADA activity only in body fluids, as shown in Figure 1.



**Fig. 1:** Combined ADA Activity in both serum and body fluid

In the suspected TB group, mean ADA values in Serum were 34.1 and 86.7 in body fluids, whereas in the control group, mean ADA serum levels were 17.3 and 42 in body fluids, as shown in Figure 2.



**Fig. 2:** Mean ADA levels in serum and Body fluids in both study groups

**Table 4:** Number of samples positive for adenosine deaminase enzyme

|          | Suspected EPTB N=50 |                 | Control N=50 |                 |
|----------|---------------------|-----------------|--------------|-----------------|
|          | Serum N(%)          | Body Fluid N(%) | Serum N(%)   | Body Fluid N(%) |
| Positive | 48(96)              | 16(32)          | 17 (34)      | 3(6)            |
| Negative | 2(4)                | 34(68)          | 33(66)       | 47(94)          |

**Table 5:** Predictive value of ADA activity in only Serum, only Body fluid,d and Combined

| Specimen type<br>Result | Serum              |                       | Body Fluids     |                       | Serum & Body Fluids |                       |
|-------------------------|--------------------|-----------------------|-----------------|-----------------------|---------------------|-----------------------|
|                         | Test group<br>(50) | Control group<br>(50) | Test group (50) | Control group<br>(50) | Test group (50)     | Control group<br>(50) |
| ADA Positive            | 34                 | 17                    | 2               | 3                     | 14                  | 0                     |
| ADA Negative            | 16                 | 33                    | 48              | 47                    | 0                   | 30                    |
| PPV                     | 66.67%             |                       | 40.00%          |                       | 100%                |                       |
| NPV                     | 67.35%             |                       | 49.47%          |                       | 100%                |                       |

### 3.1. Predictive value of the test

It was observed that both Positive Predictive value (PPV) and Negative Predictive value (NPV) for ADA detection were low when only either Serum (66.67% & 67.35%) or body fluid (40% & 49.47%) was considered for the diagnosis of EPTB. The PPV and NPV were 100% when a combined ADA activity in Serum and body fluids were considered to diagnose EPTB, as shown in Table 5.

## 4. Discussion

Tuberculosis is one of the most common infectious bacterial diseases and continues to threaten human health worldwide. Tuberculosis can be classified as pulmonary, extrapulmonary and disseminated tuberculosis. However, the mycobacterium culture of the serous fluid specimens has a relatively lower success rate; thus, its role in diagnosing extrapulmonary TB is still controversial. Due to the non-specific clinical manifestations and negative laboratory findings, it is challenging to diagnose extrapulmonary TB. Consequently, it is imperative to develop a reliable biomarker that can be used to rapidly and accurately diagnose extrapulmonary TB. The efficacy of ADA activity was studied as a biomarker for the diagnosis of extrapulmonary TB.

In our study of the group of suspected TB patients, we found that ADA activity was higher in Serum (96%) as well as body fluids (32%). Compared to the study group, the control group showed significantly low levels of ADA positivity in both Serum (32%) and body fluids (6%). These findings in our study are the first of their kind, as no literature is available for comparison.

Our present study observed that serum ADA positivity was significantly high 34(68%). ADA activity in body fluid alone was at least 2(4%). An additional eight samples were diagnosed when both Serum and ADA were considered for diagnosis. The predictive value of the test also rose to 100% when combined ADA activity was considered for the diagnosis of EPTB. These observations are the first time

observed in our study

Mean serum ADA levels were significantly higher in suspected TB patients (34.1) than in our study's control group (17.3). These observations are in concordance with other studies done by Prashant C et al., 2017<sup>16</sup> (37.12 and 15.7 respectively) and Abhijit Ninghot et al., 2019<sup>17</sup> (43.67 and 14.97 respectively)

Mean body fluid ADA levels were significantly higher in suspected TB patients (86.) than in the control group (42) in our study. Similar observations were made in other studies done by Sibel Yurt et al. 2014<sup>18</sup> (87.6 and 40.11 respectively) and Poonam Nanwani et al. 2018<sup>19</sup> (85.97 and 39.33 respectively)

## 5. Conclusion

In the present scenario of the increasing incidence of tuberculosis worldwide, particularly in developing countries, has created a need for cheaper, less time-consuming and more effective diagnostic techniques. Though X-ray and acid-fast staining are the two most common tuberculosis diagnosis methods in the developing world, these are less effective in diagnosing paucibacillary and extrapulmonary tuberculosis. Our observations suggest that serum and serosal fluid Adenosine deaminase (ADA) measurement has good prediction potential for EPTB. Hence it can be used as a supportive surrogate marker for challenging to diagnose extrapulmonary TB. ADA activity in body fluids is also a sensitive biomarker, especially when combined with serum ADA levels, and may become a routine investigation for early detection of extrapulmonary TB.

## 6. Limitations of the study

ADA may substantially reduce invasive tests like biopsy and culture; however, further studies are required at different locations and over different populations to generalize this study's results.

## 7. Source of Funding

None.

## 8. Conflicts of Interest

None.


## Acknowledgements


None.

## References

1. WHO 2012 Report: TB in South East Asia - country profile: Bangladesh. Geneva: World Health Organization; 2012.
2. Mehta JB, Dutt A, Harvill L, Mathews KM. Epidemiology of extrapulmonary tuberculosis: a comparative analysis with pre-AIDS era. *Chest*. 1991;99(5):1134–8. doi:10.1378/chest.99.5.1134.
3. Ahmed S, Fatema R, Saleh AA, Sattar H, Miah MRA. Diagnostic significance of pleural fluid adenosine Deaminase activity in tuberculous pleurisy. *Ibrahim Med Coll J*. 2011;5(1):1–5.
4. Sharma SK, Mohan A. Extrapulmonary tuberculosis. *Indian J Med Res*. 2004;120(4):316–53.
5. Bueno CE, Clemente MG, Castro BC, Martin LM, Ramos SR, Panizo AG, et al. Cytologic and bacteriologic analyzes of fluid and pleural biopsy with cop's needle. *Arch Intern Med*. 1990;150(6):1190–4.
6. Mrinal P, Subinay D. Adenosine Deaminase and its isoenzyme as a diagnostic marker in tubercular pleural effusion. *J Drug Del Ther*. 2014;4(1):18–21.
7. Sharma SK, Tahir M, Mohan A, Smith-Rohrberg D, Mishra HK, Pandey RM, et al. Diagnostic accuracy of ascitic fluid IFN-gamma and adenosine deaminase assays in the diagnosis of tuberculous ascites. *J Interferon Cytokine Res*. 2006;26(7):484–8. doi:10.1089/jir.2006.26.484.
8. Riquelme A, Calvo M, Salech F, Valderrama S, Pattillo A, Arellano M, et al. Value of adenosine deaminase (ADA) in ascitic fluid for the diagnosis of tuberculous peritonitis: a meta-analysis. *J Clin Gastroenterol*. 2006;40(8):705–10. doi:10.1097/00004836-200609000-00009.
9. Sanai FM, Bzeizi KI. Systematic review: tuberculous peritonitis—presenting features, diagnostic strategies and treatment. *Aliment Pharmacol Ther*. 2005;22(8):685–700. doi:10.1111/j.1365-2036.2005.02645.x.
10. Caws M, Wilson SM, Clough C, Drobniewski F. Role of IS6110-Targeted PCR, Culture, Biochemical, Clinical, and Immunological Criteria for Diagnosis of Tuberculous Meningitis. *Clin Microbiol*. 2000;38(9):3150–55. doi:10.1128/jcm.38.9.3150-3155.2000.
11. Mathur PC, Tiwari KK, Trikha S, Tiwari D. Diagnostic value of adenosine deaminase (ADA) activity in tubercular serositis. *Indian J Tuberc*. 2006;53:92–5.
12. Boonyagars L, Kiartiburanakul S. Use of Adenosine Deaminase for the Diagnosis of Tuberculosis: A Review. *J Infect Dis Antimicrob Agents*. 2000;27(2):111–8.
13. Song D, Lun AR, Chiu W. Diazyme adenosine Deaminase in diagnosing tuberculous pleural effusion: method evaluation and clinical experiences in a New Zealand population. *NZ J Med Lab Sci*. 2010;64(1):11–3.
14. Feres MC, Martino MC, Maldijian S, Batista F, Gabriel JA, Tufik S, et al. Laboratorial validation of an automated assay for the determination of adenosine deaminase activity in pleural fluid and cerebrospinal fluid. *J Bras Pneumol*. 2008;34(12):1033–9. doi:10.1590/s1806-37132008001200008.
15. Kashyap RS, Kainthla RP, Mudaliar AV, Purohit HJ, Taori GM, Dagainwala HF, et al. Cerebrospinal fluid adenosine deaminase activity: A complimentary tool in the early diagnosis of tuberculous meningitis. *Cerebrospinal Fluid Res*. 2006;3:5. doi:10.1186/1743-8454-3-5.
16. Chikkahonnaiah P, Jaggi S, Goyal B, Garg K, Gupta S, Jaswal S, et al. Utility of sSerumADA estimation in the diagnosis of extrapulmonary tuberculosis. *JMSCR*. 2017;5(5):21549–53.
17. Ninghot A, Mohod K, Kumar S. Evaluation of Serum Adenosine Deaminase (ADA) Values for Detection of Pulmonary and Extrapulmonary Tuberculosis. *Int J Clin Biochem Res*. 2017;4(2):106–10.
18. Yurt S, Küçükergin C, Yigitbas BA, Seçkin Ş, Tigin HC, Koşar AF, et al. Hüseyin Cem Tigin, Ayşe Filiz Koşar. Diagnostic utility of Serum and pleural levels of adenosine deaminase 1-2 and interferon- $\gamma$  in the diagnosis of pleural tuberculosis. *Multidiscip Respir Med*. 2014;9(1):12. doi:10.1186/2049-6958-9-12.
19. Nanwani P, Kapoor A, Khatri S. Diagnostic value of adenosine deaminase activity in tuberculosis & non-tuberculous lymphocytic body fluids. *J Evid Based Med Health Care*. 2018;5(4):554–61.

## Author biography

**Sateesh K**, Professor and HOD  <https://orcid.org/0000-0003-0550-4462>

**Sathya Anandam**, Associate Professor  <https://orcid.org/0000-0001-7099-7035>

**Cite this article:** Sateesh K, Anandam S. The predictive role of combined adenosine deaminase activity in Serum and body fluids to diagnose extrapulmonary tuberculosis. *IP Int J Med Microbiol Trop Dis* 2023;9(2):121-125.