

An introduction to Tropical Disease: A review article

Piyush Pujara^{1,*}, Mayur Parmar², Pratik Rupakar³, Kailash Asawa⁴, Shrikant Patel⁵

¹Senior Lecturer, ⁴Reader, Dept. of Public Health Dentistry, ⁵Senior Lecturer, Dept. of Oral Pathology, Pacific Dental College & Hospital, Rajasthan, ²Senior Lecturer, Dept. of Periodontics, Goenka Research Institute of Dental Sciences, Gujarat, ³Assistant Professor, Dept. of Dentistry, Gujarat Adani Institute of Medical Sciences, Kutch, Gujarat

***Corresponding Author:**

Email: drpiyushpujara@gmail.com

Abstract

Infectious diseases that either occur uniquely or more commonly in steamy and subtropical regions, are either more widespread in the tropics or extra tricky to prevent or control. The citizens who are the majority exaggerated by these diseases are frequently the poorest populations, whose residence is in remote, rural areas, urban slums or conflict zones. Neglected tropical diseases persevere under circumstances of scarcity and are intense approximately solely in poor populations in the developing world. The designation “tropical diseases” arise at no meticulous date and was slowly merge, as microorganisms came to be recognized as the underlying factor of diseases and had their broadcast mechanisms elucidate. In practice, the term is often taken to pass on to infectious diseases that flourish in burning, moist circumstances, such as malaria, leishmaniasis, schistosomiasis, onchocerciasis, lymphatic filariasis, Chagas disease, African trypanosomiasis, and dengue. A number of the organisms that grounds tropical diseases are bacteria and viruses, conditions that may be recognizable to the majority people as these types of organisms’ grounds sickness common. Fewer well recognized are those more compound organisms usually referred to as parasites. Global warming donates in introducing and scattering more diseases internationally, but in short term; it defiantly would not reason any epidemic circumstances by itself.

Keywords: Global warming, Malaria, Tropical diseases, WHO

Access this article online
Website: www.innovativepublication.com
DOI: 10.5958/2455-6807.2016.00011.8

Introduction

During current history, tropical areas of the world were more harshly exaggerated by infectious diseases in contrast to the mild world. Main reasons why infectious diseases can flourish in such regions can be establish in together environmental and biological factors that hold up high levels of biodiversity of pathogens, vectors and hosts, but also in social factors that weaken pains to manage these diseases.¹ Such infectious diseases are recognized just as tropical diseases and tropical medicine has come out as an significant regulation for their study.^{2,3}

A few of these diseases are reasons by protozoa, such as malaria, leishmaniasis, Chagas’ disease and sleeping sickness while others are due to worms, including chistosomiasis, onchocerciasis and lymphatic filariasis. Viral disease is dengue fever. The eight WHO tropical diseases are broadcasts to humans by various means, but forever comprise a vector that is usually a hematophagous insect.

The designation “tropical diseases” was not invented by the WHO and has been part of the medical vocabulary since the 19th century. It arose at no particular date and was gradually consolidated, as

microorganisms came to be acknowledged as the causal factors of diseases and had their transmission mechanisms elucidated.^{4,5}

There is surely a burly part of underdevelopment in tropical diseases, a delayed result from colonial times, but there is also a certain tropical destiny, the effect of heterogeneous biological and biological evolution. The diseases that humanity obtain all through history from its associates in the voyage– such as dogs, cats, rodents, birds and even our close up relatives.

Different example of Tropical Disease

Malaria is a good preliminary summit for our analysis. The microorganism’s that because the diseases are parasitic protozoa of red blood cells; encompass four species of the genus Plasmodium. Among every the Earth’s animals, these four species parasitize only humans, to whom they are broadcasts by mosquitoes that transport them from the ill to the fit. The four type of plasmodia have escorts human race since Homo sapiens and Plasmodium ssp left Africa jointly to inhabit the Earth. In every curve of the planet – cold, temperate or hot this duo has relied on the involvement of a priory accessible species of Anopheles to proliferate and perpetuate it. With the exemption of the glacial regions, no occupied nook or cranny supervised to shun malaria, which becomes a worldwide scourge. Similarly, no nation – current or very old – deals with to escape.

Though, malaria is listed by the WHO among the tropical diseases and, indeed, that is how it is seen today. If we examine the current world distribution of

malaria, we will see that it prevails and has greater incidence in countries located between the Tropic of Cancer and the Tropic of Capricorn, that is, between latitudes 27° 23' north and south. Exceptions are the Middle Eastern countries, notably Afghanistan, northern India and parts of southern China, where malaria is left over. However, it is factual that malaria is these days rigorous in the tropics and this information carry us to an early conclusion: if malaria was one time pandemic, the disease was not and surely is not at the present connected to any tropical biogeographic fate. In fact, this cause is under growth and the resulting scarcity of the tropical populations. Conscious of this information, agencies such as Unicef, the World Bank and the WHO itself start on the particular Program for Research and Training in Tropical Diseases(TD R) thirty years ago, alert on infectious diseases that excessively trouble the “deprived and exile populations” of the earth.⁶

Tuberculosis corresponds to the most important reason of demise linked with infectious diseases internationally, and its occurrence is on the increase in tropical areas owing to the communication between tuberculosis and HIV epidemics. In numerous regions of the earth this disease mainly affects youthful adults, and rising fight of the pathogen to antimicrobial drugs is a troublesome sign.⁵

Diarrhea leftovers the majority widespread diseases that also influence children under 5 years of age, leads to a significant death in early days. Rotavirus remnants the mainly frequent reason of severe diarrhea sickness, although a countless of countries have seen severe epidemics owing to the numerous resistant bacterium *Shigelladysenteriae* that grounds dysentery. Leishmaniasis is a cluster of diseases due to parasite of the genus *Leishmania*, which is endemic in 88 countries and leads to major morbidity and mortality. Strongyloidiasis a frequent reason of disease in tropical and subtropical areas, as humid climate is appropriate for parasite endurance. The clinical appearance varies with the standing of the host's immune system, and the infection can be classified as acute, chronic and severe.

Onchocerciasis, too recognized as river blindness, is an illness caused by the filarial nematode, *Onchocerca volvulus*. Larvae of this worm can shift beneath the skin and go through the eye, then ensuing in sight injury and blindness.^{6,7}

Lymphatic filariases are evenly limited to the tropics. These are caused by the nematode worms *Wuchereriabancrofti*, originate all above the tropical world, and *Brugyamalayi*, originate merely in south-east Asia. The worms can turn out to be fairly abundant after consecutive infections and can then occlude the lymphatic vessels, foremost to the buildup of lymph upstream. Filariases have been accounts in the upper Nile since antiquity and the figurine of at least one pharaoh shows signs of elephantiasis of the leg. Extremely inflated scrota are ordinary together in

African statuettes 1,500 years aged and in modern patients.⁸

It is expected that further than 1 billion people are at danger of constricting one of the filariae, which are broadcasts from human to human by the omnipresent and cosmopolitan ordinary mosquitoes. Dengue fever's *Aedes* and malaria's *Anopheles* can also put out filariae, but these insects are not the vital ones. Debatably, the most significant of all is the frequent mosquito of the genus *Culex*. The agents used for treatment of filariasis are the same as the ones for onchocerciasis, are donated free of charge by the laboratories that produce them – a rare but dignifying example of the pharmaceutical industry.⁹

Management and prevention actions

Tropical diseases can be put under control by distributing medications, improving living conditions and building waste sanitation facilities. Furthermore, planned programs that enable nutritional support to those who live in high-risk areas can help to strengthen resistance of a human organism to these diseases. Moreover, community-wide vaccination programs can create a distinction, and some vaccines for abandoned tropical diseases have entered the clinical pipeline. Improved collaboration, namely among the public sector and vaccine producers in developing countries, could furthermore endorse information distribution and cost savings.

Shortage in water and sanitation absolutely donate to the serious disease load imposed by infectious diarrhea, hepatitis, intestinal nematodes like hookworm and leishmaniasis, thus it should also be addressed.¹⁰

Repellents should obtain more importance as a vital public health measure in tropical regions where vectors bite in the premature sunset or when confronted with epidemics. Bed nets should be also utilized to decrease nocturnal broadcast of this disease.

Conclusion

Global warming contributes in introducing and spreading more diseases globally, but in short term, it defiantly would not cause any epidemic situations by itself.

References

1. <http://www.nejm.org/doi/pdf/10.1056/nejmra064142>. Accessed on 21/08/2015.
2. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3438653/>. Accessed on 22/09/2015.
3. <http://www.prb.org/pdf10/neglectedtropicaldiseases.pdf>. Accessed on 12/08/2015.
4. http://www.who.int/neglected_diseases/diseases/NTD_Report_APPMG.pdf. Accessed on 11/01/2016.
5. Meunier YA, Hole M, Shumba T, Swanner BJ. Tropical Diseases: A Practical Guide for Medical Practitioners and Students. OUP USA, 2013; pp. 1-106.
6. COX, F. E. G. History of human parasitology. *Clinical Microbiology Reviews*, v.15, p. 595-612, 2002.

7. BACELLAR, R. C. *Brazil's contribution to tropical medicine and malaria*. Rio de Janeiro: Gráfica Olímpica Editora, 1963.
8. FENWICK, A. et al. Implementation of human schistosomiasis control: challenges and prospects. *Advances in Parasitology*, v.61, p. 567-622, 2006.
9. FERREIRA, L. F. et al. Paleoparasitology of Chagas' disease revealed by infected tissues from Chilean mummies. *Acta Tropica*, v.75, p. 79-84, 2000.
10. Shetty NP, Shetty PS. Epidemiology of Disease in the Tropics. In: Cook GC, Zumla A, editors. *Manson's Tropical Diseases*. Elsevier Health Sciences, 2009; pp. 19-34.

How to cite this article: Pujara P, Parmar M, Rupakar P, Asawa K, Patel S. An introduction to Tropical Disease: A review article. *International Journal of Medical Microbiology and Tropical Diseases* 2016;2(3):81-83.