

## Bacteriological profile and its antibiotic susceptibility in patients with Urinary Tract Infection at Tertiary Care Hospital, Valsad, Gujarat

Vicky P. Gandhi<sup>1,\*</sup>, Mehul Patel<sup>2</sup>, Alka Nerurkar<sup>3</sup>

<sup>1</sup>Assistant Professor, <sup>2</sup>Tutor, <sup>3</sup>Professor & HOD, Dept. of Microbiology,

**\*Corresponding Author:**

Email: vickygandhi15@gmail.com

### Abstract

**Introduction:** Urinary tract infection is one of the most common bacterial infections seen in humans and major cause of morbidity. The causative agents for urinary tract infection vary from place to place and they also vary in their sensitivity and resistance pattern with the widespread availability of antimicrobial drugs. UTI is difficult to treat because of appearance of pathogens with increasing resistant pattern to antimicrobial drugs.

**Aim:** This study is carried out to know the bacteriological profile and its antibiotics sensitivity pattern in patients with UTI at tertiary care hospital, Valsad.

**Materials and Method:** Total 771 mid-stream urine samples were collected from suspected UTI patients. They were tested for bacterial culture and all Culture positive samples were tested for antimicrobial susceptibility by Kirby- Bauer disc diffusion method.

**Results:** Out of total 771 samples, isolates were detected in 302(39.16%) samples. Incidence of infection was more common in females (55%) as compared to males (45%). *Escherichia coli* (36.75%) was the most common pathogens followed by *Klebsiella* spp. (18.21%), Gram positive organisms (16.22%), *Pseudomonas* spp. (12.25%) and *Candida* spp.(8.60%). Isolated pathogenic organism shows high resistance to commonly use antibiotic and sensitive to higher generation of fluoroquinolones and carbapenemase.

**Conclusion:** In this study, females were mostly affected and most common organisms were *E.coli* and *Klebsiella*. The commonly isolated pathogens have been changing and increasing resistance pattern is observed due to indiscriminate use of antibiotics and alerting us to update effective empirical treatment regularly.

**Keywords:** Urinary tract infection, Antimicrobial susceptibility, Resistance pattern

### Introduction

Urinary tract infections (UTI) are the most common infectious diseases in clinical practice.<sup>(1)</sup> This problem spans all age groups, beginning from neonates to the geriatrics age group.<sup>(2,3)</sup> UTI represent the second most common microbial infection after respiratory tract infections, encountered in medical practice.<sup>(4)</sup> It was estimated that in a year UTI was the cause of 100,000 cases of hospitalizations, 1 million visits to the emergency department and 7 million visits to outpatient department all over the world.<sup>(5,6)</sup>

UTI is defined as the presence of growth of more than 10<sup>5</sup> colony forming unit (CFU) of bacteria per ml of urine for asymptomatic individual and 10<sup>3</sup> for symptomatic individual.<sup>(7)</sup> It is estimated that the incidence is greater in women as compare to men due to anatomical predisposition, large bacterial load in urogenital mucosa, sexual activity and pregnancy. Infection in men is uncommon through the 5<sup>th</sup> decade of life, after that the incidence of infection is high because of enlargement of prostate interferes with emptying of the bladder.<sup>(7,8)</sup>

The most common pathogenic organisms of UTI are *Escherichia coli*, *staphylococcus saprophyticus* and less common pathogens are *Klebsiella* spp., *Proteus* spp., *Pseudomonas* spp., *Enterococci* spp, and *Candida albicans*.<sup>(9)</sup> Treatment of UTI cases is often started empirically and therapy is based on information

determined from the antimicrobial resistance pattern of the urinary pathogens. UTI caused by bacteria have been showing increasing trends even though there is availability and use of antibiotics.<sup>(10)</sup>

The extensive and inappropriate use of antimicrobial agents has invariably resulted in the development of antibiotic resistance which in recent years, has become a major problem worldwide.<sup>(11)</sup> To ensure appropriate treatment, knowledge of the organisms that cause infection and its antimicrobial sensitivity pattern is mandatory.<sup>(11,12)</sup>

### Materials and Method

The study was conducted at Microbiology department, GMERS medical college and hospital, Valsad, Gujarat from January 2014 to December 2015. Total 771 mid stream urine samples were collected from clinically suspected UTI patients. Urine samples were cultured on Nutrient agar, Sheep blood agar by semi-quantitative method and on MacConkey agar.<sup>(13,14)</sup> All plates were incubated at 37°C aerobically for 24 hours. The plates were examined macroscopically and microscopically for bacterial growth. A growth of  $\geq 10^5$  colony forming units/ml consider as significant bacteriuria, according to the standard Kass criteria.<sup>(13,15)</sup>

Bacterial pathogens were identified by routine gram reactions, motility testing and biochemical reaction as per Standard Microbiology techniques.<sup>(13,15)</sup>

Antimicrobial susceptibility testing was performed by Kirby-Bauer disc diffusion method, using commercially available discs on Muller- Hinton agar.<sup>(15,16)</sup>

## Result


The study was conducted from January 2014 to December 2015 at tertiary care hospital Valsad. In study duration, total 771 urine samples were processed, out of which 302 (39.16%) samples shows growth of pathogenic organism. The most common isolated was *E.coli* in 111(36.75%) samples, which was followed by *Klebseilla spp.* 55 (18.21%), *Pseudomonas spp.* 37 (12.25%), *Candida spp.*26 (8.60%), *Enterococcus spp.*19 (6.29%) and *S.aureus* 13 (4.30%). (Table 1) The prevalence rate in female 166 (55%) is more as compare to male 136(45%).

**Table 1: Frequency of different isolated organism**

Organism	Frequency of isolates in numbers (Percentage)
<i>E.coli</i>	111 (36.75%)
<i>Klebseilla spp.</i>	55 (18.21%)
<i>Pseudomonas spp.</i>	37 (12.25%)
<i>Candida spp.</i>	26 (8.60%)
<i>Enterococcus spp.</i>	19 (6.29%)
Coagulase negative <i>staphylococcus</i>	16 (5.29%)
<i>S.aureus</i>	13 (4.30%)
<i>Acinetobacter spp.</i>	11 (3.64%)
<i>Proteus spp.</i>	7 (2.31%)
<i>Providentia spp.</i>	2 (0.66%)
Other organism*	5 (1.65%)


\*includes *citrobacter spp.*, *morgenella spp.*, *S.maltophilia* and *streptococcus spp.*

In antibiotic susceptibility testing, *E.coli* was highly sensitive to Imipenam (83.78%), Nitrofurantoin (81.08%) and Levofloxacin (72.27%). *E.coli* was shows resistance to amikacin (38.73%), gentamycin (44.14%) and piperacillin-tazobactum (45%).




**Fig. 1: Antibiotic sensitivity of *Escherichia coli***

The antibiotic pattern of *Klebseilla spp* shows high sensitivity to levofloxacin, imipenam, amikacin, gentamycin and piperacillin-tazobactum. *Klebseilla spp.* shows high resistance to ampicilline, cefaclor, cefotaxime.(Fig. 2) *Pseudomonas spp.* species were highly sensitive to piperacillin-tazobactum, imipenam, amikacin. (Fig. 3)



**Fig. 2: Antibiotic sensitivity of *Klebsiella* species**



**Fig. 3: Antibiotic sensitivity of *Pseudomonas* species**

In gram positive organism sensitivity, all organisms *S.aureus*, *Enterococcus* species and coagulase negative *staphylococcus* are sensitive for vancomycin (100%) and linezolid(100%).(Table 2)

**Table 2: Antibiotic sensitivity patterns of gram positive organism**

Antimicrobial agents	Sensitivity of organism in percentage		
	<i>S.aureus</i>	<i>Enterococcus spp.</i>	CONS
Tetracycline	92.30	63.15	81.25
Oxacilline	38.46	21.05	68.75
Azithromycin	30.76	00.00	53.84
Clindamycin	84.61	00.00	81.25
Cotrimoxazole	30.76	00.00	31.25
Chloramphenicol	61.53	47.36	75
Vancomycin	100	100	100
Levofloxacin	76.92	68.42	92.30
Linezolid	100	100	100

## Discussion

Urinary tract infection causes major burden in health care services because of high prevalence in community and hospitals by different pathogenic organisms. It also contributes to increasing antimicrobial drugs resistance both at community and hospital level. So continues surveillance of antibiotic sensitivity is necessary at local level for better management of patient by choosing appropriate

antimicrobial therapy and to decrease antibiotic drug resistance.

In our study, prevalence of uropathogens was 39.16%, which was near similar (39.6%) to study done by Nilofar S et al at Anand district, Gujarat.<sup>(17)</sup> some other studies in India shows prevalence rate from 31-38%.<sup>(18,19)</sup>

The most common isolate found in our study was *E.coli* (36.75%), which was followed by *Klebsiella* spp. (18.21%) and *Pseudomonas* spp. (12.25%). Our study's

result was comparable to several studies done in India like study done by Dnyaneshwari Ghadage et al, at Pune in 2016 shows *E.coli* isolate rate was 41.3% and *Klebsiella* spp. was 18.5%.<sup>(20)</sup> Other studies also shows *E.coli*, *Klebsiella* spp. and *pseudomonas* spp. as most common isolated pathogenic organism.<sup>(5,17)</sup>

In our study, Gram negative organisms show high resistance to ampicilline, cotrimoxazole and 3<sup>rd</sup> and 4<sup>th</sup> generation of cephalosporine. Gram negative organisms were more sensitive to imipenam, nitrofurantoin and levofloxacin which are similar as study done in India.<sup>(5,21)</sup> There is a high resistance developed in ampicilline and cotrimoxazole which were previously used as treatment of choice in last decades but indiscriminate and empirical use of these drugs in every patient resulted in high resistance in these drugs.

Gram positive cocci show 100% sensitivity to vancomycin and linezolid and resistance to azithromycin, cotrimoxazole and oxacilline which are similar to studies done in India.<sup>(22,23)</sup>

### Conclusion

Our study concludes *E. coli* as most common organism causing urinary tract infection with female predominance. There is increasing resistance to commonly used antibiotics due to indiscriminate use of antibiotics without culture and sensitivity report. Periodic antimicrobial sensitivity surveillance is necessary for formulating antibiotic policy and based on antibiotic policy start earlier and effective empirical treatment for better management and reduces morbidity to patients.

### References

- Alonto AM, Urinary tract infections. Mahon CR, Lehman DC, Textbook of Diagnostic microbiology, 3<sup>rd</sup> edition, Elsevier, 2007.
- Kalpana S, Hegadi SS, Ramesh K. Characterization and [2] antimicrobial susceptibility testing of uropathogens from urinary tract infections. *Int J Curr Microbiol Appl Sci.* 2015;4(2):1010-16.
- Raju CB, Urinary tract infection – A suitable approach, journal Indian academy of clinical medicine 2001;2(4):331-334.
- Hryniewicz K, Szczypa K et al, Antibiotic susceptibility of bacterial strains isolated from urinary tract infections in Poland. *J Antimicrobial chemotherapy* 2001;47(6):773-780.
- Razak SK, Gurushantappa V. Bacteriology of urinary tract infection and antibiotic susceptibility pattern in a tertiary care hospital in South India. *Int J Med Sci Public Health.* 2012;1(2):109-12.
- Patel S, Taviad PP, Sinha M, Javadekar TB, Chaudhari VP. Urinary tract infections (UTI) among patients at G.G. Hospital & Medical College, Jamnagar. *Natl J Community Med.* 2012;3(1):138-41.
- Bailey & Scott's, Diagnostic Microbiology, 12<sup>th</sup> edition, Infection of urinary tract, Page No.842-855.
- Sohail M, Khurshid M, Characteristics and antibiotic resistance of urinary tract pathogens isolated from Punjab, Pakistan. *Jundishapur journal of microbiology* 2015;8(7):01-04.
- Salek SB, 1992, Infective syndrome in medical microbiology, 4<sup>th</sup> edition, pp 740.
- Wilson ML, Gaido L, Laboratory diagnosis of urinary tract infection in adult patients. *Clin Infect Dis,* 2004;38:1150-1158.
- Goldstein FE, Antibiotic susceptibility of bacterial strains isolated from patients with community acquired urinary tract infection in France. *Eur J Clin Microbiol Infect Ds* 2000;19:112-117.
- Ashkenazi S, Samra Z, Uropathogens of various childhood populations and their antibiotics susceptibility. *Pedia Infec Ds* 1991;10:742-746.
- Winn WC, Allen SD, Koneman EW et al, Koneman's colour atlas and textbook of diagnostic microbiology, 6<sup>th</sup> edition. Williams and Wilkins 2006.
- Kass E, Bacteriuria and diagnosis of urinary tract. *Arch Internal medicine,* 100: 709-714, 1957.5.
- Collee JG, Frase AG, Marmion BP, et al. Laboratory strategy in the diagnosis of infective syndrome. Mackie and McCartney Practical Medical Microbiology. 14<sup>th</sup> edition 2006.
- Bauer AW, Kirbey WMM, Sherris JC, et al. Antibiotic susceptibility testing by a standardised single disc method. *Am J Clin Patho* 1966;45:493-496.
- Nilofar S, Trivedi S et al. Microbiological profile and antibiotic susceptibility pattern of uropathogens in Anand district, Gujarat. *Journal of Biomedical and Pharmaceutical research,* 4:4, July-august 2015;49-56.
- Patel S, Taviad PKP et al. Urinary tract infection among patients at GG hospital and medical college, Jamnagar, Gujarat. *National J Community Medicine* 2012;3(1):138-141.
- Ritu Aggrawal, Maneesh Goyal. Bacterial isolates and their antibiotic sensitivity profile recovered from urine samples in NCR, Ghaziabad *JEMDS* 2014;3(28):7831-7836.
- Dnyaneshwari Ghadage et.al. Bacteriological profile and anti-biogram of urinary tract infections at tertiary care hospital, Pune. *National journal of laboratory medicine* 2016 Oct Vol5(4):20-24.
- Bashir MF, Qazi JI et al. Diversity of urinary tract pathogens and drug resistant isolates group of Pakistanis. *Tropical journal of Pharmaceutical Research* Sept. 2008; 7(3):1025-1031.
- Sood S, Gupta R. Antibiotic resistance pattern of community acquired uropathogens at a tertiary care hospital in Jaipur, Rajasthan. *Indian journal community medicine* 2012;37(1):39-44.
- Laxminarayana SA, Chavan Sk et al. Bacterial pathogens in urinary tract infection and antibiotic susceptibility pattern from a teaching hospital, Bengaluru, India. *International J Curr Microbiol Applied Science* 2015;4(11):731-36.